

PPH Risk Assessment Management for 3rd Stage

Score less than 6	Score 6 - 9	Score 10 or more
Standard 3rd stage management	Follow Green Action Plus	Follow Yellow action PLUS:
Syntometrine 1ML IM	Iv access - 1 x 18G cannula	IV access - 2 18G IV cannulas
(Ergometrine 500 microg with oxytocin 5 units)	Group and hold and FBE	
(give antiemetic unless declines)	Collected and sent	
	Inform the O&G Registrar	
OR if BP $\geq$ 140/90 then use		Consider :
Oxytocin (Syntocinon) 10 units IM or		Crossmatching 2 units of blood
5 units IV (if IV gives slow over 3 - 5 min)		Inform the O&G consultant and Anaesthetic Registrar

Antenatal PPH Risk Assessment

Placenta Previa/Accreta	10
Antepartum haemorrhage recurrent or >50mls	10
Multiple Pregnancy	6
Current Hb < 85	6
Parity >6	6
Pre-eclampsia/Gestational hypertension (with or without Medication)	4
Maternal Platelet Disorder	3
Previous PPH or Retained Placenta	3
Macrosomia > 4500g	3
Maternal Iron Deficiency	3
Polyhydramnios	3
Current BMI > 35	2
Uterine Fibroids/Uterine anomaly/Intrauterine Fetal Demise	2
Recurrent APH	2

Intrapartum PPH Risk Assessment

Augmentation of Labour	2
Chorioamnionitis or Sepsis/Pyrexia in Labour	2
Prolonged 1st stage of labour > 12 hours (active)	2
> 12 hours of Syntocinon	2
Prolonged 2nd stage of labour > 2 hours	2